

Figure 12. *Neophylloceras hetonaiense* Matsumoto, 1942a from the Etanpakk Formation. **A–D**, HMG-1703 from Loc. 2; **E–H**, HMG-1704 from Loc. 2; **I–L**, HMG-1706 from Loc. 14; **M–O**, HMG-1707 from a float calcareous concretion found near Loc. 14; **P–R**, HMG-1702 from a float calcareous concretion found near Loc. 1; **S**, HMG-1705 from a float calcareous concretion found near Loc. 8.

Hypophylloceras (Neophylloceras) hetonaiense Matsumoto.
Morozumi, 1985, p. 15, pl. 1, figs. 1–5.

Neophylloceras nera (Forbes). Matsumoto and Toshimitsu,
1996, p. 3, pls. 1, 2.

Neophylloceras cf. nera (Forbes). Maeda *et al.*, 2005, p. 55,
fig. 24.1–24.4.

Neophylloceras sp. Shigeta *et al.*, 2015, p. 110, fig. 3A–L.

Lectotype.—Specimen designated by Matsumoto
(1959, p. 5) is GK. H3801a. This is the original of

Matsumoto (1942, p. 675) from the sandy mudstone of Unit IVb of the Hakobuchi Formation at loc. H12b (Matsumoto, 1942b) in the Hobetsu area, Hokkaido.

Material examined.—One specimen, HMG-1702, from a float calcareous concretion found near Loc. 1; two specimens, HMG-1703, 1704, from Loc. 2; one specimen, HMG-1705, from a float calcareous concretion found near Loc. 8; one specimen, HMG-1706, from Loc. 14; one specimen, HMG-1707, from a float calcareous concretion found near Loc. 14.

Description.—Very involute, very compressed shell with elliptical whorl-section, arched venter, rounded ventral shoulders and slightly convex flanks with maximum whorl width at mid-flank. Umbilicus very narrow and deep with moderately high, vertical wall and rounded shoulders. Ornamentation consists of fine, dense, weak flexuous lirae, which arise at umbilical seam, sweep gently forward across inner flank, and then strengthen and become rectiradiate at mid-flank before passing straight across venter. Lirae gradually develop into slightly more distant, narrowly raised ribs, which increase in strength as diameter increases. Broad undulations sometimes appear on inner flank. Suture line consists of numerous deeply incised elements with phylloid terminals.

Measurements (mm):

Specimen no.	D	U	H	W	U/D	W/H
HMG-1704	23.0	1.9	13.2	7.9	0.08	0.60
HMG-1703	24.8	1.9	14.0	8.0	0.08	0.57
HMG-1706	31.1	2.4	17.1	10.3	0.08	0.60
HMG-1702	39.0	3.1	21.8	11.7	0.08	0.54
HMG-1707	43.1	3.5	24.5	13.0	0.08	0.53

Remarks.—Matsumoto and Toshimitsu (1996) assigned three specimens from the lower Maastrichtian (Unit IVc of the Hakobuchi Formation) in the Hobetsu area to *Neophylloceras nera* (Forbes, 1846), but these specimens are identical to *N. hetonaiense* in having very weak flexuous lirae or ribs. *Neophylloceras nera* is characterized by sigmoidal lirae on its inner flanks (Kennedy and Henderson, 1992, p. 389, pl. 1, figs. 10–12, pl. 15, figs. 1, 2). Juvenile specimens reported as *N. cf. nera* by Maeda et al. (2005) from the Maastrichtian in the Makarov area, southern Sakhalin and *Neophylloceras* sp. by Shigeta et al. (2015) from the upper Maastrichtian in the Akkeshi Bay area, eastern Hokkaido should probably be assigned to *N.*

hetonaiense, because they have very weak flexuous lirae. *Neophylloceras lambertense* Usher, 1952 from Vancouver Island, British Columbia was earlier regarded as a synonym of *N. hetonaiense* by Jones (1963).

Juvenile specimens reported as *Neophylloceras hetonaiense* from the Campanian in northern Hokkaido by Matsumoto (1984a) and Matsumoto and Miyauchi (1984) should probably be assigned to *N. ramosum* (Meek, 1857), because juveniles of both species are sometimes very close and larger specimens referable to *N. hetonaiense* have never been collected from northern Hokkaido.

Occurrence.—*Neophylloceras hetonaiense* is known from the upper Campanian in Antarctica (Spath, 1953), the Maastrichtian in Southwest Japan (Matsumoto and Morozumi, 1980; Morozumi, 1985), Hobetsu and Akkeshi Bay areas in Hokkaido (Matsumoto, 1942a; Shigeta et al., 2015), southern Sakhalin (Zonova et al., 1993), southern Alaska (Jones, 1963), Vancouver Island (Usher, 1952) and California (Matsumoto, 1959).

Suborder Ammonitina Hyatt, 1889

Superfamily Desmoceratoidea Zittel, 1895

Family Pachydiscidae Spath, 1922

Genus *Pachydiscus* Zittel, 1884

Type species.—*Ammonites neubergicus* Hauer, 1858.

Pachydiscus sp.

Figure 13A–C

Material examined.—One specimen, HMG-1708, from Loc. 12.

Description.—Moderately involute, fairly compressed shell with elliptical whorl-section, arched venter, rounded ventral shoulders, and slightly convex flanks with maximum whorl width at mid-flank. Fairly narrow umbilicus with moderately high, nearly vertical wall and rounded umbilical shoulders. Ornamentation consists of radial primary ribs that begin at elongate umbilical bullae and intercalated secondary rib that begin on the umbilical shoulder. Both sets of ribs pass over venter in a broad convex arch.

Remarks.—The described specimen is very similar




Figure 13. *Pachydiscus*, *Anagaudryceras* and *Gaudryceras* from the Etanpakk Formation. **A–C**, *Pachydiscus* sp., HMG-1708 from Loc. 12; **D–G**, *Anagaudryceras matsumotoi* Morozumi, 1985, HMG-1709 from Loc. 1; **H–J**, *Gaudryceras izumiense* Matsumoto and Morozumi, 1980; H, I, HMG-1712 from Loc. 2; J, HMG-1714 from Loc. 12.

to the juvenile shells of *Pachydiscus kamishakensis* Jones, 1963 and *P. hazzardi* Jones, 1963 from the Maastrichtian of southern Alaska, but a definitive assignment cannot be made.

Suborder Lytoceratina Hyatt, 1889
Superfamily Tetragonitoidea Hyatt, 1900
Family Gaudryceratidae Spath, 1927
Genus *Anagaudryceras* Shimizu, 1934

Type species.—*Ammonites sacya* Forbes, 1846.

Anagaudryceras matsumotoi Morozumi, 1985

Figure 13D–G

Anagaudryceras matsumotoi Morozumi, 1985, p. 29, pl. 9, fig. 1, text-fig. 7; Matsumoto, 1985, p. 27, pl. 4, figs. 1–10; Matsumoto, 1988, p. 183, pl. 51, fig. 3; Ando *et al.*, 2001, pl. 1, figs. 12–14; Maeda *et al.*, 2005, p. 81, fig. 39.1–39.15; Shigeta *et al.*, 2015, p. 112, figure 5A–P.

Zelandites varuna (Forbes). Zonova *et al.*, 1993, p. 148, pl. 98, fig. 4; Yazykova, 1994, p. 289, pl. 1, fig. 8.

Holotype.—GK. H6882, figured by Morozumi (1985, p. 29, pl. 9, fig. 1, text-fig. 7), from the Maastrichtian *Pachydiscus* aff. *subcompressus* Zone in the Shimonada Formation of the Izumi Group on Awaji Island, Southwest Japan.

Material examined.—One specimen, HMG-1709,




Figure 14. *Gaudryceras izumiense* Matsumoto and Morozumi, 1980 from the Etanpakk Formation. **A–C**, HMG-1716 from Loc. 14; **D–F**, HMG-1710 from Loc. 1.

from Loc. 1.

Description.—Fairly evolute shell with whorl nearly as high as broad. Whorl-section circular with arched venter, indistinct ventral shoulders and gently convex flanks. Maximum whorl width occurs at a

slight distance below mid-flank. Umbilicus moderately wide with moderately high, rounded umbilical wall. Ornamentation consists of very fine slightly sinuous growth lines, which pass straight across venter.

Measurements (mm):



Figure 15. *Gaudryceras izumiense* Matsumoto and Morozumi, 1980 from the Etanpakk Formation. **A, B**, HMG-1717 from a float calcareous concretion found near Loc. 14; **C, D**, HMG-1718 from a float calcareous concretion found near Loc. 14; **E**, HMG-1710 from Loc. 1.

Specimen no.	D	U	H	W	U/D	W/H
HMG-1709	29.0	10.8	11.6	12.0	0.37	1.03

Remarks.—Zonova *et al.* (1993, pl. 98, fig. 4) and Yazykova (1994, pl. 1, fig. 8) assigned a specimen from the Maastrichtian in the Pugachevo area, southern

Sakhalin, to *Zelandites varuna*, but as Maeda *et al.* (2005) and Shigeta *et al.* (2015) earlier pointed out, the specimen is identical to *Anagaudryceras matsumotoi* with respect to whorl-section, mode of coiling and ornamentation.



Figure 16. *Gaudryceras izumiense* Matsumoto and Morozumi, 1980 from the Etanpakk Formation. **A, B**, HMG-1711 from Loc. 2; **C, D**, HMG-1715 from a float calcareous concretion found near Loc. 12; **E, F**, HMG-1713 from Loc. 9.

Occurrence.—*Anagaudryceras matsumotoi* is known from the Maastrichtian of southern Sakhalin (Matsumoto, 1988; Zonova *et al.*, 1993; Maeda *et al.*, 2005), northern Hokkaido (Matsumoto, 1985), eastern Hokkaido (Matsumoto, 1985; Shigeta *et al.*, 2015) and Southwest Japan (Morozumi, 1985).

Genus ***Gaudryceras*** Grossouvre, 1894

Type species.—*Ammonites mitis* Hauer, 1866.

Gaudryceras izumiense Matsumoto and Morozumi,

1980

Figures 13H–J, 14–16

Gaudryceras tenuiliratum Yabe, Kobayashi, 1931, p. 639, pl. 10.

Gaudryceras izumiense Matsumoto and Morozumi, 1980, p. 12, pl. 11, fig. 1, pl. 12, fig. 1, pl. 13, fig. 1; Shigeta *et al.*, 2010, p. 205, figs. 2–5.

Gaudryceras venustum Matsumoto, 1984c, p. 5, pl. 3, figs. 1, 2; Matsumoto and Toshimitsu, 1995, p. 2, pls. 1–8.

Holotype.—OMNH.M1125, figured by Matsumoto and Morozumi (1980, p. 13, pl. 11, fig. 1), from the Maastrichtian Azenotani Formation of the Izumi Group at a roadside cliff about 800 m west of Sobura (Loc. 8), Kaizuka, Osaka Prefecture, Southwest Japan.

Material examined.—One specimen, HMG-1710, from Loc. 1; two specimens, HMG-1711, 1712, from Loc. 2; one specimen, HMG-1713, from Loc. 9; one specimen, HMG-1714, from Loc. 12; one specimen, HMG-1715, from a float calcareous concretion found near Loc. 12; one specimen, HMG-1716, from Loc. 14; two specimens, HMG-1717, 1718, from float calcareous concretions found near Loc. 14.

Description.—Early to middle growth stages (up to 80 mm in diameter): Very evolute, slightly depressed shell with arched venter, indistinct ventral shoulders, and slightly convex flanks with maximum whorl width at mid-flank. Umbilicus wide with moderately high, vertical wall and rounded shoulders. Ornamentation consists of very fine, dense, slightly sinuous lirae, which arise at umbilical seam and pass over venter in a broad convex arch. Intercalation of lirae occurs on umbilical shoulder and lower flank. Each whorl has variable dense or distant, rounded, collar-like ribs, running parallel to lirae; and each rib is immediately followed by a shallow constriction.

Later growth stage (over 80 mm in diameter): As shell grows larger, whorl section becomes more compressed. Lirae gradually develop into slightly more distant, narrowly raised ribs, which increase in strength as diameter increase. On body chamber of mature shell, ribs become much coarser and distant, and collar-like ribs become more frequent.

Measurements (mm):

Specimen no.	D	U	H	W	U/D	W/H
HMG-1715	47.0	26.1	14.2	15.3	0.56	1.08
HMG-1711	52.1	28.0	13.9	15.0	0.54	1.08

HMG-1710	99.0	52.0	37.0	35.0	0.53	0.95
HMG-1716	115.0	48.2	39.1	36.2	0.42	0.93
HMG-1718	173.0	58.5	68.0	—	0.34	—
HMG-1717	181.0	56.0	70.0	—	0.31	—
HMG-1710	220.0	66.0	88.2	74.1	0.30	0.84

Remarks.—Shigeta *et al.* (2010) considered *Gaudryceras venustum* to be conspecific with *G. izumiense* because specimen HMG-1541 from the Hobetsu area exhibits ornamentation characteristic of *G. izumiense* on one side, while the other side is abraded and appears very similar to *G. venustum*.

Occurrence.—*Gaudryceras izumiense* is known from the lower Maastrichtian in Southwest Japan (Matsumoto and Morozumi, 1980), the Hobetsu area in Hokkaido (Shigeta *et al.*, 2010) and southern Alaska (Shigeta *et al.*, 2010).

Family Tetragonitidae Hyatt, 1900

Genus **Tetragonites** Kossmat, 1895

Type species.—*Ammonites timotheanus* Pictet, 1847.

Tetragonites terminus Shigeta, 1989

Figure 17

Tetragonites terminus Shigeta, 1989, p. 338, figs. 11E, F, 13.8–13.10; Ando and Ando, 2002, pl. 1, figs. 7, 8.

Holotype.—UMUT MM18635-1, figured by Shigeta (1989, p. 338, fig. 13.8), from the sandy mudstone of Unit IVb of the Hakobuchi Formation at loc. H12d (Matsumoto, 1942b) in the Hobetsu area, Hokkaido.

Material examined.—Two specimens, HMG-1719, 1720, from Loc. 1; two specimens, HMG-1721, 1722, from Loc. 6; two specimens, HMG-1723, 1724, from Loc. 12; one specimen, HMG-1725, from Loc. 13, one specimen, HMG-1726, from Loc. 14.

Description.—Fairly involute, fairly depressed shell with sub-rectangular whorl section, broadly rounded venter, rounded ventral shoulders, and slightly convex flanks with maximum whorl width near umbilical shoulder. Umbilicus fairly narrow with moderately high, nearly vertical wall and rounded shoulders. Ornamentation consists only of distant constrictions and very fine, growth lines, which are prorsiradiate on flanks, but become slightly sinuous at ventral shoulders before crossing venter in a very shallow concave arch.




Figure 17. *Tetragonites terminus* Shigeta, 1989 from the Etanpakk Formation. **A–D**, HMG-1719 from Loc. 1; **E, F**, HMG-1721 from Loc. 6; **G, H**, HMG-1722 from Loc. 6; **I–L**, HMG-1720 from Loc. 1; **M–P**, HMG-1723 from Loc. 12; **Q–T**, HMG-1724 from Loc. 12; **U–X**, HMG-1726 from Loc. 14; **Y–AB**, HMG-1725 from Loc. 13.

Measurements (mm):

Specimen no.	D	U	H	W	U/D	W/H
HMG-1725	13.1	3.1	5.9	6.5	0.24	1.10
HMG-1724	16.0	3.8	6.7	8.4	0.24	1.26
HMG-1722	19.0	4.6	8.3	9.2	0.24	1.11
HMG-1719	19.6	4.8	8.2	8.2	0.24	1.00
HMG-1720	22.4	5.5	9.0	9.3	0.25	1.03
HMG-1726	22.6	5.6	10.7	11.2	0.25	1.05

Remarks.—Shigeta (1989) studied the early internal shell morphology of *Tetragonites terminus* and revealed that the initial chamber and ammonitella sizes range from 0.925 to 1.050 mm and 1.700 to 1.900 mm in diameter respectively, which are unusually large for Mesozoic ammonoids (Landman *et al.*, 1996; De Baets *et al.*, 2015). Such large initial chambers and ammonitellas are partly visible in HMG-1719, 1720, 1722, 1724, 1725 and 1726.

Occurrence.—*Tetragonites terminus* is known from the lower Maastrichtian *Nostoceras hetonaiense* Zone in the Hobetsu area, Hokkaido (Shigeta, 1989).

Tetragonites popetensis Yabe, 1903

Figures 18–21

Tetragonites popetensis Yabe, 1903, p. 48, pl. 7, figs. 4, 6; Matsumoto and Miyauchi, 1984, p. 52, pl. 23, fig. 3; Matsumoto, 1988, p. 178, pl. 50, figs. 3, 4; Zonova *et al.*, 1993, p. 155, pl. 209, fig. 2; Yazykova, 1994, p. 293, pl. 3, fig. 1; Naruse *et al.*, 2000, fig. 3.4; Ando and Ando, 2002, pl. 2, figs. 4, 5; Maeda *et al.*, 2005, p. 88, figs. 38.9–38.11, 38.14, 38.15, 42.5–42.11, 43, 44; Kurihara and Kano, 2006, pl. 3, fig. 2; Takahashi *et al.*, 2007, pl. 1, fig. 4; Misaki and Maeda, 2009, fig. 8E, F; Shigeta *et al.*, 2016, p. 333, figs. 7D–H, 8D, E, 24A, B.

Epigonceras epigonus (Kossmat). Usher, 1952, pl. 2, figs. 6, 7, pl. 3, fig. 1.

Lytoceras (Tetragonites) henleyense Anderson, 1958, p. 185, pl. 12, fig. 5, pl. 41, fig. 7.

Tetragonites superstes (Hoepen). Matsumoto and Miyauchi, 1984, p. 52, pl. 23, fig. 2.

Tetragonites glabrus (Jimbo). Shigeta, 1989, p. 334, fig. 12.4–12.7.

Saghaliinites maclerei (White). Haggart, 1989, p. 186, pl. 8.1, figs. 7–11.

Saghaliinites cala (Forbes). Zonova *et al.*, 1993, p. 155, pl. 209, figs. 3, 4.




Figure 18. *Tetragonites popetensis* Yabe, 1903 from the Etanpakk Formation. **A–D**, HMG-1730 from Loc. 2; **E–H**, HMG-1732 from Loc. 6; **I–L**, HMG-1735 from Loc. 12; **M, N**, HMG-1733 from Loc. 12; **O–R**, HMG-1738 from Loc. 14; **S–V**, HMG-1728 from a float calcareous concretion found near Loc. 1.

Holotype.—UNUT MM7460, figured by Yabe (1903, p. 48, pl. 7, fig. 4), from the Upper Cretaceous of the Sanushube (Sanushibe) River area in the Hobetsu area, Hokkaido.

Material examined.—One specimen, HMG-1727, from Loc. 1; two specimens, HMG-1728, 1729, from float calcareous concretions found near Loc. 1; two specimens, HMG-1730, 1731, from Loc. 2; one specimen, HMG-1732, from Loc. 6; one specimen,




Figure 19. *Tetragonites popetensis* Yabe, 1903 from the Etanpakk Formation. **A, B**, HMG-1736 from a float calcareous concretion found near Loc. 12; **C**, HMG-1731 from Loc. 2; **D, E**, HMG-1729 from a float calcareous concretion found near Loc. 1.



Figure 20. *Tetragonites popetensis* Yabe, 1903 from the Etanpakk Formation. **A–C**, HMG-1727 from Loc. 1; **D, E**, HMG-1729 from a float calcareous concretion found near Loc. 1.




Figure 21. *Tetragonites popetensis* Yabe, 1903 from the Etanpakk Formation. **A–C**, HMG-1734 from a float calcareous concretion found near Loc. 11; **D–F**, HMG-1737 from Loc. 13.

HMG-1733, from Loc. 11; one specimen, HMG-1734, from a float calcareous concretion found near Loc. 11; one specimen, HMG-1735, from Loc. 12; one specimen, HMG-1736, from a float calcareous concretion found near Loc. 12; one specimen, HMG-1737, from Loc. 13; one specimen, HMG-1738, from Loc. 14.

Description.—Moderately evolute, fairly depressed shell (early growth stages) to fairly compressed shell (later growth stages) with sub-quadratus whorl section, broadly rounded venter, rounded ventral shoulders, and slightly convex flanks with maximum whorl width near umbilical shoulder. Umbilicus moderately wide with moderately high, nearly vertical wall and rounded shoulders. Ornamentation consists only of very fine, growth lines, which are prorsiradiate on flanks, but become slightly sinuous at ventral shoulders before crossing venter in a shallow concave arch. Specimen HMG-1729 (Figure 19D) exhibits a conspicuous constriction near its aperture.

Measurements (mm):

Specimen no.	D	U	H	W	U/D	W/H
HMG-1730	11.7	3.9	4.7	5.0	0.33	1.06
HMG-1733	18.6	5.7	30.6	31.2	0.28	1.02
HMG-1732	20.1	6.5	7.9	8.3	0.31	1.05
HMG-1735	22.7	7.0	9.6	10.5	0.31	1.09
HMG-1738	30.0	9.4	12.4	14.0	0.31	1.13
HMG-1734	61.2	19.0	26.2	28.2	0.31	1.08
HMG-1737	63.0	19.1	27.0	28.6	0.30	1.06
HMG-1728	71.0	19.9	30.6	31.2	0.28	1.02
HMG-1729	127.4	42.8	49.0	48.2	0.34	0.98

Remarks.—Shigeta (1989) demonstrated that *Tetragonites glabrus* of Turonian to Campanian age, exhibits a remarkably wide variation in shell form and regarded *T. popetensis* as a synonym of *T. glabrus*. However, Shigeta's specimens, which exhibit a small umbilicus that he assigned to *T. glabrus* from sample AW1001A, should actually be attributed to *Pseudophyllites indra* (Forbes, 1846). Because Shigeta's sample (AW1001A) was heterogeneous and consisted of specimens of both *Pseudophyllites indra* (Forbes, 1846) and *T. popetensis*. *Tetragonites popetensis* should be regarded as an independent species as earlier pointed out by Maeda *et al.* (2005, p. 88).

Certain specimens assigned to *Saghalinites* by Haggart (1989, pl. 8.1, figs. 7–11) and Zonova *et al.*, (1993, pl. 209, figs. 3, 4) as well as a specimen described as *Tetragonites superstes* by Matsumoto and Miyauchi (1984, pl. 23, fig. 2) and specimens described as *Epigonoceras epigonum* by Usher (1952, pl. 2, figs. 6, 7, pl. 3, fig. 1) are identical to *Tetragonites popetensis* in having a moderately evolute shell with a sub-quadratus whorl section and should be synonymized with *T. popetensis*. The holotype of *Lytoceras (Tetragonites) henleyense* exhibits a similar shell shape and ornamentation as *T. popetensis*, and is probably conspecific. However, it is necessary to study a large number of specimens in order to define the variation in shell form of *L. (T.) henleyense*.

All specimens of *Tetragonites popetensis* reported from the Campanian are less than 70 mm in diameter (Shigeta, 1989). A mature shell, which is characterized by septal approximation of the last two septa, was reported from the Campanian in the Cape Soya area by Matsumoto and Miyauchi (1984, pl. 23, fig. 3) and its diameter is over 65 mm. In contrast, the largest specimen of *T. popetensis* from the Soya Hill area (HMG-1729) is 127.4 mm in diameter, which is nearly twice as large as the Campanian mature specimens. This evidence suggests that the mature shell size of *T. popetensis* increased significantly during the Maastrichtian.

Occurrence.—*Tetragonites popetensis* is abundant from the Santonian to the lower Maastrichtian in Hokkaido and Sakhalin (Shigeta, 1989; Maeda *et al.*, 2005; Shigeta *et al.*, 2016) and from the middle Campanian in Wakayama, southwestern Japan (Misaki and Maeda, 2009). The species is also known from the Coniacian to Campanian in California, Washington and British Columbia (Usher, 1952; Anderson, 1958; Haggart, 1989).

Suborder Ancyloceratina Wiedmann, 1966

Superfamily Turrilitoidea Gill, 1871

Family Diplomoceratidae Spath, 1926

Genus *Diplomoceras* Hyatt, 1900

Type species.—*Baculites cylindracea* Defrance, 1816.

Diplomoceras sp.

Figure 22A–E

Material examined.—One specimen, HMG-1739, from Loc. 7; one specimen, HMG-1740, from a float calcareous concretion found near Loc. 8.

Description.—HMG-1740, a curved portion of phragmocone, with elliptical whorl section ($H = 20.2$ mm, $W = 18.0$ mm, $W/H = 0.89$). HMG-1739 is a body chamber fragment. Ornamentation consists of numerous, regularly spaced, straight ribs.

Remarks.—The fragmental nature of the specimens preclude a definitive assignment.

Family Nostoceratidae Hyatt, 1894

Genus **Nostoceras** Hyatt, 1894

Type species.—*Nostoceras santoni* Hyatt, 1894.

Nostoceras sp.

Figure 22F, G

Material examined.—One specimen, HMG-1741, from a float calcareous concretion found near Loc. 11.

Description.—HMG-1741 consists of fragments of a phragmocone, probably the last portion of sinistrally helical coiling, and a body chamber. Whorl cross section nearly circular for phragmocone ($H = 24.1$ mm, $W = 23.3$ mm, $W/H = 0.97$) and elliptical for body chamber ($H = \text{ca. } 49$ mm). Shell surface ornamented with dense, oblique ribs and two rows of tubercles, which appear rather irregularly on every second to fourth rib. As shell grows, ribs increase in strength and tubercles become stronger.

Remarks.—The described specimen is somewhat similar to *Nostoceras hornbyense* (Whiteaves, 1895), but the fragmental nature precludes a definitive species assignment.

Family Baculitidae Gill, 1871

Genus **Baculites** Lamarck, 1799

Type species.—*Baculites vertebralis* Lamarck, 1801.

Baculites regina Obata and Matsumoto, 1963

Figures 23, 24

Baculites regina Obata and Matsumoto, 1963, p. 85, pl. 22, figs. 3–6, pl. 23, figs. 1, 2, pl. 24, figs. 1–5, pl. 25, figs. 3–5, pl. 27, figs. 1, 6, 7, 9, text-figs. 191–196, 200–214; Matsumoto and Morozumi, 1980, p. 24; Klinger and Kennedy, 2001, p. 192.

Holotype.—UNUT MM7716b, figured by Obata and Matsumoto (1963, p. 85, pl. 24, fig. 1, text-fig. 210), from the lower Maastrichtian Azenotani Shale Member of the Izumi Group at Kuratani, Shinke, Sennan, Osaka Prefecture, Southwest Japan.

Material examined.—One specimen, HMG-1742, from Loc. 1; one specimen, HMG-1743, from Loc. 6; five specimens, HMG-1744–1748, from float calcareous concretions found near Loc. 8; five specimens, HMG-1749–1753, from Loc. 11; one specimen, HMG-1754, from a float calcareous concretion found near Loc. 11.

Description.—Moderately tapered, straight or gently arched shell. Whorl section rounded subtrigonal, with narrowly rounded venter, indistinct ventral shoulders, gently convex flanks, flattened or broadly rounded dorsum. As shell grows in length, whorl section becomes more compressed. Shell surface almost smooth in early growth stage, but as shell grows, it develops weak to strong crescentic nodes on flanks and numerous, fine, oblique ribs on ventral half of flank.

Measurements (mm):

Specimen no.	<i>H</i>	<i>W</i>	<i>W/H</i>
HMG-1745	5.9	4.8	0.81
HMG-1750	7.6	6.0	0.79
HMG-1751	7.6	5.9	0.78
HMG-1749	8.6	6.5	0.76
HMG-1754	8.8	6.6	0.75
HMG-1752	9.3	6.8	0.73
HMG-1746	10.8	7.3	0.68
HMG-1748	10.8	7.9	0.72
HMG-1753	13.3	9.2	0.69
HMG-1747	14.3	10.1	0.70
HMG-1742	14.4	10.0	0.69
HMG-1744	19.7	11.0	0.56

Remarks.—*Baculites regina* closely resembles *B. subanceps* Haughton, 1925 in having crescentic nodes on its flanks, but differs by having numerous, fine, oblique ribs on the ventral half of its flank. *Baculites subanceps* has a corrugated venter and oval whorl section.

Occurrence.—*Baculites regina* is known from the Lower Maastrichtian Azenotani Shale Member of the




Figure 22. *Diplomoceras* and *Nostoceras* from the Etanpakk Formation. A–E, *Diplomoceras* sp.; A–D, HMG-1740 from a float calcareous concretion found near Loc. 8; E, HMG-1739 from Loc. 7; F, G, *Nostoceras* sp., HMG-1741 from a float calcareous concretion found near Loc. 11.




Figure 23. *Baculites regina* Obata and Matsumoto, 1963 from the Etanpakk Formation. **A–D**, HMG-1743 from Loc. 6; **E–I**, HMG-1742 from Loc. 1; **J–X**, specimens from a float calcareous concretion found near Loc. 8; **J–N**, HMG-1745; **O–S**, HMG-1746; **T–X**, HMG-1744; **Y–AH**, specimens from a float calcareous concretion found near Loc. 8; **Y–AC**, HMG-1747; **AD–AH**, HMG-1748.




Figure 24. *Baculites regina* Obata and Matsumoto, 1963 from the Etanpakk Formation. A–E, HMG-1754 from a float calcareous concretion found near Loc. 11; F–AD, specimens from a float calcareous concretion found near Loc. 11; F–J, HMG-1749; K–O, HMG-1750; P–T, HMG-1751; U–Y, HMG-1752; Z–AD, HMG-1753.

Izumi Group at Sennan, Osaka Prefecture, Southwest Japan.

Conclusions

1. Zircon geochronology reveals that the ages of the tuffs from the lowest and uppermost parts of the Etanpakku Formation in the Soya Hill area are 72.6 ± 1.6 Ma and 70.6 ± 1.2 Ma, respectively, which infer a late Campanian to earliest middle Maastrichtian age.

2. An early Maastrichtian (latest Cretaceous) ammonoid fauna is reported from the middle part of the Etanpakku Formation of the Yezo Group exposed along the Onishibetsu and Sarukotsu rivers in the Soya Hill area, northernmost Hokkaido, northern Japan.

3. The fauna comprises nine species belonging to eight genera: *Neophylloceras hetonaiense*, *Tetragonites popetensis*, *T. terminus*, *Gaudryceras izumiense*, *Anagaudryceras matsumotoi*, *Pachydiscus* sp., *Baculites regina*, *Diplomoceras* sp. and *Nostoceras* sp.

4. The fauna correlates with the lower part of the *Gaudryceras izumiense* Zone of the upper lower Maastrichtian in the Izumi Group in Southwest Japan.

5. The *Gaudryceras izumiense* Zone in Hokkaido suggests that similar faunas may have existed in the North Pacific realm during late early Maastrichtian time.

Acknowledgments

We are very grateful to Noboru Sasaki (Sapporo, Hokkaido) and the Hokkaido Regional Forest Office for their co-operation in the field. We thank Jim Jenks (West Jordan, Utah) for his helpful suggestions and improvement of the English text. This study was financially supported by grants from the National Museum of Nature and Science in 2016 and JSPS KAKENHI Grant Number JP25400502 to Y. S.

References

- Anderson, F. M., 1958: Upper Cretaceous of the Pacific Coast. *Geological Society of America Memoir*, vol. 71, p. 1–378.
- Ando, H., Tomosugi, T. and Kanakubo, T., 2001: Upper Cretaceous to Paleocene Hakobuchi Group, Nakatonbetsu area, northern Hokkaido—lithostratigraphy and megafossil biostratigraphy. *Journal of the Geological Society of Japan*, vol. 107, p. 142–162. (in Japanese with English abstract)
- Ando, M. and Ando, H., 2002: Depositional facies and megafossil biostratigraphy of the Upper Cretaceous Hakobuchi Group in the Soya Hill area, northern Hokkaido. *Bulletin of the Nakagawa Museum of Natural History*, vol. 5, p. 1–21. (in Japanese with English abstract)
- Arkell, W., 1950: A classification of the Jurassic ammonites. *Journal of Paleontology*, vol. 24, p. 354–364.
- Arkell, W. J., 1957: Introduction to Mesozoic Ammonoidea. In, Arkell, W. J., Furnish, W. M., Kummel, B., Miller, A. K., Moore, R. C., Schindewolf, O. H., Sylvester-Bradley, P. C. and Wright, C. W. eds., *Treatise on Invertebrate Paleontology, Part L, Mollusca 4, Cephalopoda, Ammonoidea*, p. L81–129. Geological Society of America, New York and University of Kansas Press, Lawrence.
- Black, L. P., Kamo, S. L., Allen, C. M., Davis, D. W., Aleinikoff, J. N., Valley, J. W., Mundil, R., Campbell, I. H., Korsch, R. J., Williams, I. S. and Foudoulis, C., 2004: Improved $^{206}\text{Pb}/^{238}\text{U}$ microprobe geochronology by the monitoring of a trace-element-related matrix effect; SHRIMP, ID-TIMS, ELA-ICP-MS and oxygen isotope documentation for a series of zircon standards. *Chemical Geology*, vol. 205, p. 115–140.
- Corfu, F., Hanchar, J. M., Hoskin, P. W. O. and Kinny, P., 2003: An atlas of zircon textures. In, Hanchar, J. M. and Hoskin, P. W. O. eds., *Zircon: Reviews in Mineralogy and Geochemistry*, p. 278–286. Mineralogical Society of America, Washington DC.
- De Baets, K., Landman, N. H. and Tanabe, K., 2015: Ammonoid embryonic development. In, Klug, C., Korn D., De Baets, K., Kruta I. and Mapes, R. H. eds., *Ammonoid Paleobiology: From Anatomy to Ecology*, p. 113–205. Topics in Geobiology, vol. 43, Springer, Dordrecht, Heidelberg, New York and London.
- Defrance, M. J. L., 1816: *Dictionnaire des Sciences naturelles, dans lequel on traite méthodiquement des différents Etres de la Nature* 3, 492 p. Levrault, Paris and Strasbourg.
- Forbes, E., 1846: Report on the fossil Invertebrata from southern India, collected by Mr. Kaye and Mr. Cunliffe. *Transactions of the Geological Society of London, Series 2*, vol. 7, p. 97–174.
- Gill, T., 1871: Arrangement of the families of mollusks. *Smithsonian Miscellaneous Collections*, vol. 227, p. 1–49.
- Grossouvre, A. de., 1894: Recherches sur la Craie supérieure. Deuxième partie: paléontologie. Les ammonites de la Craie supérieure. *Mémoires du Service de la Carte Géologique Détailée de la France*, p. 1–264.
- Haggart, J. W., 1989: New and revised ammonites from the Upper Cretaceous Nanaimo Group of British Columbia and Washington State. *Geological Survey of Canada Bulletin* 396, p. 181–221.
- Hauer, F. R. von., 1858: Ueber die Cephalopoden der Gosauschichten. *Beitrag zur Paläontographie von Österreich*, Band 1, p. 7–14.
- Hauer, F. R. von., 1866: Neue Cephalopoden aus den Gosaugebilden der Alpen. *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften in Wien*,

- mathematisch-Naturwissenschaftliche*, Band 53, p. 300–308.
- Haughton, S. H., 1925: Notes on some Cretaceous fossils from Angola (Cephalopoda and Echinoidea). *Annals of the South African Museum*, vol. 22, p. 263–288.
- Hoskin, P. W. and Black, L. P., 2000: Metamorphic zircon formation by solid-state recrystallization of protolith igneous zircon. *Journal of Metamorphic Geology*, vol. 18, p. 423–439.
- Hyatt, A., 1889: Genesis of the Arietidae. *Smithsonian Contributions to Knowledge*, no. 673, p. 1–238.
- Hyatt, A., 1894: Phylogeny of an acquired characteristics. *Proceedings of the American Philosophical Society*, vol. 32, p. 349–647.
- Hyatt, A., 1900: Cephalopoda. In, Zittel K. A. ed., *Textbook of Palaeontology*, English ed., translated by C. R. Eastman, p. 502–592. Macmillan, London and New York.
- Jimbo, K., 1894: Beiträge zur Kenntniss der Fauna der Kreideformation von Hokkaido. *Palaeontologische Abhandlungen Neue Folge*, Band 2, p. 1–48.
- Jones, D., 1963: Upper Cretaceous (Campanian and Maastrichtian) ammonites from southern Alaska. *United States Geological Survey Professional Paper*, no. 432, p. 1–53.
- Kennedy, W. J. and Henderson, R. A., 1992: Non-heteromorph ammonites from the upper Maastrichtian of Pondicherry, South India. *Palaeontology*, vol. 35, p. 381–442.
- Kinny, P. D., Wijbrans, J. R., Froude, D. O., Williams, I. S. and Compston, W., 1990: Age constraints on the geological evolution of the Narreyer Gneiss Complex, Western Australia. *Australian Journal of Earth Sciences*, vol. 37, p. 51–69.
- Klinger, H. C. and Kennedy, W. J., 2001: Stratigraphic and geographic distribution, phylogenetic trends and general comments on the ammonite family Baculitidae Gill, 1871 (with an annotated list of species referred to the family). *Annals of the South African Museum*, vol. 107, p. 1–290.
- Kobayashi, T., 1931: On the Izumi Sandstone Series in the Izumi mountain range. *Journal of the Geological Society of Tokyo*, vol. 38, p. 629–640.
- Kossmat, F., 1895: Untersuchungen über die Südindische Kreideformation. Teil 1. *Beiträge zur Paläontologie und Geologie Österreich-Ungarns und des Orients*, Band 9, p. 97–203.
- Kurihara, K. and Kano, M., 2006: Minoru Yamashita's collection donated to the Mikasa City Museum. *Bulletin of the Mikasa City Museum*, no. 10, p. 1–35. (in Japanese with English abstract)
- Lamarck, J. B. P. A. de M. de, 1799: Prodrome d'une nouvelle classification des coquilles. *Mémoires de la Société d'Histoire naturelle de Paris*, p. 63–90.
- Lamarck, J. B. P. A. de M. de, 1801: *Système des Animaux sans vertebrae*, 432 p. Deterville, Paris.
- Landman, N. H., Tanabe, K. and Shigeta, Y., 1996: Ammonoid embryonic development. In, Landman, N. H., Tanabe, K. and Davis, R. A. eds., *Ammonoid Paleobiology*, p. 343–405. Topics in Geobiology, vol. 13, Plenum Press, New York and London.
- Ludwig, K. R., 2003: *User's manual for Isoplot 3.00. A geochronological toolkit for Microsoft Excel*. Berkeley Geochronology Center Special Publication No. 4, 70 p. Berkeley Geochronology Center, Berkeley.
- Maeda, H., Shigeta, Y., Fernando, A. G. S. and Okada, H., 2005: Stratigraphy and fossil assemblages of the Upper Cretaceous System in the Makarov area, southern Sakhalin, Russian Far East. *National Science Museum Monographs*, no. 31, p. 25–120.
- Matsumoto [= Matumoto], T., 1942a: A short note on the Japanese Cretaceous Phylloceratidae. *Proceedings of the Imperial Academy, Tokyo*, vol. 18, p. 674–676.
- Matsumoto [= Matumoto], T., 1942b: Fundamentals in the Cretaceous stratigraphy of Japan, Part 1. *Memoirs of the Faculty of Science, Kyushu Imperial University, Series D, Geology*, vol. 1, p. 129–280.
- Matsumoto, T., 1954: *The Cretaceous System in the Japanese Islands*, 324 p. Japan Society for the Promotion of Science, Tokyo.
- Matsumoto, T., 1959: Upper Cretaceous ammonites of California. Part 2. *Memoirs of the Faculty of Science, Kyushu University, Series D, Geology*, Special vol. 1, p. 1–172.
- Matsumoto, T., 1977: Some heteromorph ammonites from the Cretaceous of Hokkaido. *Memoirs of the Faculty of Science, Kyushu University, Series D, Geology*, vol. 23, p. 303–366.
- Matsumoto, T., 1979: Palaeontological descriptions Part 1. Some new species of *Pachydiscus* from the Tombetsu and the Hobetsu valleys. *Memoirs of the Faculty of Science, Kyushu University, Series D, Geology*, vol. 24, p. 50–64.
- Matsumoto, T., 1984a: Ammonites from the Upper Campanian of the Teshio Mountains. *Palaeontological Society of Japan, Special Paper*, no. 27, p. 5–32.
- Matsumoto, T., 1984b: Concluding remarks. *Palaeontological Society of Japan, Special Paper*, no. 27, p. 77–91.
- Matsumoto, T., 1984c: Some gaudryceratid ammonites from the Campanian and Maastrichtian of Hokkaido Part 1. *Science Report of the Yokosuka City Museum*, no. 32, p. 1–10.
- Matsumoto, T., 1985: Three species of *Anagaudryceras* from the Campanian and Maastrichtian of Hokkaido. *Science Report of the Yokosuka City Museum*, no. 33, p. 22–29.
- Matsumoto, T., 1988: Notes on some Cretaceous ammonites from South Sakhalin held at Tohoku University, Sendai. *Science Report of the Tohoku University, 2nd series (Geology)*, vol. 59, p. 177–190.
- Matsumoto, T. and Miyauchi, T., 1984: Some Campanian ammonites from the Soya area. *Palaeontological Society of Japan, Special Paper*, no. 27, p. 33–76.
- Matsumoto, T. and Morozumi, Y., 1980: Late Cretaceous ammonites from the Izumi Mountains, Southwest Japan. *Bulletin of the Osaka Museum of Natural History*, no. 33, p. 1–31.
- Matsumoto, T. and Ohara, J., 1971: Relations of Cretaceous and Tertiary strata in the Soya area, Hokkaido. *Science Reports of the Department of Geology, Kyushu University*, vol. 11, p. 17–34. (in Japanese with English abstract)
- Matsumoto, T. and Toshimitsu, S., 1992: On a leading ammonite species *Pachydiscus kobayashii* from the Hobetsu district, Hokkaido. *Bulletin of the Hobetsu*

- Museum*, no. 8, p. 1–16.
- Matsumoto, T. and Toshimitsu, S., 1995: Restudy of *Gaudryceras venustum* Matsumoto from the Hobetsu district, Hokkaido. *Bulletin of the Hobetsu Museum*, no. 11, p. 1–16.
- Matsumoto, T. and Toshimitsu, S., 1996: A phylloceratid ammonite species from the Maastrichtian of the Hobetsu district, Hokkaido. *Bulletin of the Hobetsu Museum*, no. 12, p. 1–8.
- Matsumoto, T., Toshimitsu, S. and Noda, M., 1993: On a Maastrichtian (Cretaceous) inoceramid species *Sphenoceramus hetonaianus* (Matsumoto) from the Hobetsu distinct, Hokkaido. *Bulletin of the Hobetsu Museum*, no. 9, p. 1–20.
- Matsushita, K., Mitani, K., Ishiyama, S. and Osanai, H., 1964: *Explanatory Text of the Geological Map of Japan, Scale 1:50000, Onishibetsu (Asahikawa-10)*, 31 p. Geological Survey of Hokkaido, Sapporo (in Japanese with English abstract).
- Meek, F. B., 1857: Descriptions of new organic remains from the Cretaceous rocks of Vancouver's Island. *Transactions of the Albany Institute*, vol. 4, p. 37–49.
- Misaki, A. and Maeda, H., 2009: Lithostratigraphy and biostratigraphy of the Campanian–Maastrichtian Toyajo Formation in Wakayama, southwestern Japan. *Cretaceous Research*, vol. 30, p. 1398–1414.
- Morozumi, Y., 1985: Late Cretaceous (Campanian and Maastrichtian) ammonites from Awaji Island, Southwest Japan. *Bulletin of the Osaka Museum of Natural History*, no. 39, p. 1–58.
- Murphy, M. A. and Rodda, P. U., 2006: California Early Cretaceous Phylloceratidae (Ammonoidea). *University of California, Riverside, Campus Museum Contribution*, vol. 7, p. 1–97.
- Naruse, H., Maeda, H. and Shigeta, Y., 2000: Newly discovered Late Cretaceous molluscan fossils and inferred K/T boundary in the Nemuro Group, eastern Hokkaido, northern Japan. *Journal of the Geological Society of Japan*, vol. 106, p. 161–164. (in Japanese with English abstract)
- Obata, I. and Matsumoto, T., 1963: A monograph of the Baculitidae from Japan. Part 2. *Memoirs of the Faculty of Science, Kyushu University, Series D, Geology*, vol. 13, p. 77–92.
- Osanai, H., Mitani, K. and Kitagawa, Y., 1959: *Explanatory Text of the Geological Map of Japan, Scale 1:50000, Soya and Soya-misaki (Asahikawa-4, 1)*, 52 p. Geological Survey of Hokkaido, Sapporo (in Japanese with English abstract).
- Osanai, H., Mitani, K., Ishiyama, S. and Matsushita, K., 1963: *Explanatory Text of the Geological Map of Japan, Scale 1:50000, Nakatonbetsu (Asahikawa-21)*, 58 p. Geological Survey of Hokkaido, Sapporo (in Japanese with English abstract).
- Pictet, F. J., 1847: Description des mollusques fossiles qui se trouvent dans le Grès Vers des environs de Genève. *Mémoires de la Société de Physique et d'Histoire Naturelle de Genève*, vol. 11, p. 257–412.
- Salfeld, H., 1924: *Die Bedeutung der Konservativstämme für die Stammesentwicklung der Ammonoideen*, 16 p. Leipzig.
- Schiøtte, L., Compston, W. and Bridgwater, D., 1988: Late Archaean ages for the deposition of clastic sediments belonging to the Malene supracrustals, southern West Greenland: evidence from an ion probe U–Pb zircon study. *Earth and Planetary Science Letters*, vol. 87, p. 45–58.
- Shigeta, Y., 1989: Systematics of the ammonite genus *Tetragonites* from the Upper Cretaceous of Hokkaido. *Transactions and Proceedings of the Palaeontological Society of Japan, New Series*, no. 156, p. 319–342.
- Shigeta, Y., Tanabe, K. and Izukura, M., 2010: *Gaudryceras izumiense* Matsumoto and Morozumi, a Maastrichtian ammonoid from Hokkaido and Alaska and its biostratigraphic implications. *Paleontological Research*, vol. 14, p. 202–211.
- Shigeta, Y., Nishimura, T. and Nifuku, K., 2015: Middle and late Maastrichtian (latest Cretaceous) ammonoids from the Akkeshi Bay area, eastern Hokkaido, northern Japan and their biostratigraphic implications. *Paleontological Research*, vol. 19, p. 107–127.
- Shigeta, Y., Izukura, M., Nishimura, T. and Tsutsumi, Y., 2016: Middle and late Campanian (late Cretaceous) ammonoids from the Urakawa area, Hokkaido, northern Japan. *Paleontological Research*, vol. 20, p. 322–366.
- Shimizu, S., 1934: Ammonites. In, Shimizu S. and Obata T. eds., *Cephalopoda*, p. 1–137. Iwanami's lecture series of Geology and Palaeontology, Tokyo. (in Japanese)
- Shimizu, S., 1935: The Upper Cretaceous cephalopods of Japan. Part 1. *Journal of the Shanghai Science Institute, Section 2*, vol. 1, p. 159–226.
- Spath, L. F., 1922: On the Senonian ammonite fauna of Pondoland. *Transactions of the Royal Society of South Africa*, vol. 10, p. 113–148.
- Spath, L. F., 1926: On new ammonites from the English Chalk. *Geological Magazine*, vol. 63, p. 77–83.
- Spath, L. F., 1927: Revision of the Jurassic cephalopod fauna of Kachch (Cutch), part 1. *Memoirs of the Geological Survey of India, Palaeontologia Indica, New series*, vol. 9, memoir 2, p. 1–71.
- Spath, L. F., 1953: The Upper Cretaceous cephalopod fauna of Graham Land. *Falkland Islands Dependencies Survey, Scientific reports*, no. 3, p. 1–60.
- Stacey, J. S. and Kramers, J. D., 1975: Approximation of terrestrial lead isotope evolution by a two-stage model. *Earth and Planetary Science Letters*, vol. 26, p. 207–221.
- Suess, E., 1865: Über Ammoniten. *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften in Wien, mathematisch-Naturwissenschaftliche*, vol. 52, p. 71–89.
- Takahashi, K. and Ishiyama, S., 1968: *Explanatory Text of the Geological Map of Japan, Scale 1:50000, Numakawa (Asahikawa-9)*, 46 p. Geological Survey of Hokkaido, Sapporo (in Japanese with English abstract).
- Takahashi, A., Hikida, Y., Jenkins, R. G. and Tanabe, K., 2007: Stratigraphy and megafauna of the Upper Cretaceous Yezo Supergroup in the Teshionakagawa area, northern Hokkaido, Japan. *Bulletin of the Mikasa City Museum*, no. 11, p. 25–59.
- Tsutsumi, Y., Horie, K., Sano, T., Miyawaki, R., Momma, K., Matsubara, S., Shigeoka, M. and Yokoyama, K., 2012: LA-ICP-MS and SHRIMP ages of zircons in chevkinite and monazite tuffs from the Boso

- Peninsula, Central Japan. *Bulletin of the National Museum of Nature and Science, Series C*, vol. 38, p. 15–32.
- Usher, J. L., 1952: Ammonite faunas of the Upper Cretaceous rocks of Vancouver Island, British Columbia. *Geological Survey of Canada Bulletin* 21, p. 1–182.
- Whiteaves, J. F., 1895: Notes on some fossils from the Cretaceous rocks of British Columbia, with descriptions of two species that appear to be new. *Canadian Record of Science*, vol. 6, p. 313–318.
- Wiedmann, J., 1966: Stammesgeschichte und System der posttriadischen Ammonoideen, ein Überblick (2. Teil). *Neues Jahrbuch fuer Geologie und Palaeontologie, Abhandlungen*, Band 127, p. 13–81.
- Williams, I. S., 1998: U–Th–Pb geochronology by ion microprobe. In, McKibben M. A., Shanks W. C. P. and Ridley W. I. eds., *Applications of Microanalytical Techniques to Understanding Mineralizing Processes*, p. 1–35. Reviews in Economic Geology 7, Society of Economic Geologists, Littleton.
- Williams, I. S. and Claesson, S., 1987: Isotopic evidence for the Precambrian provenance and Caledonian metamorphism of high grade paragneisses from the Seve Nappes, Scandinavian Caledonides. *Contributions to Mineralogy and Petrology*, vol. 97, p. 205–217.
- Yabe, H., 1903: Cretaceous Cephalopoda from the Hokkaido. Part 1: *Lytoceras*, *Gaudryceras*, and *Tetragonites*. *Journal of the College of Science, Imperial University of Tokyo*, vol. 18, p. 1–55.
- Yazykova [= Yazikova], E. A., 1994: Maastrichtian ammonites and biostratigraphy of the Sakhalin and the Shikotan Islands, Far East Russia. *Acta Geologica Polonica*, vol. 44, p. 277–303.
- Zittel, K. A., 1884: Cephalopoda. In, Zittel, K. A. ed., *Handbuch der Palaeontologie, Band 1, Abt. 2, Lief 3*, p. 329–522. Oldenbourg, Munich and Leipzig.
- Zittel, K. A., 1895: *Grundzüge der Palaeontologie*, 971 p. Oldenbourg, Munich and Leipzig.
- Zonova, T. D., Kazintsova, L. I. and Yazykova, E. A., 1993: *Atlas of Index Fossils in the Cretaceous Fauna of Sakhalin*, 327 p. Nedra, St. Petersburg. (in Russian; original title translated)

Yasunari Shigeta, Masataka Izukura, and Yukiyasu Tsutsumi, 2017. An early Maastrichtian (latest Cretaceous) ammonoid fauna from the Soya Hill area, Hokkaido, northern Japan. *The Bulletin of the Hobetsu Museum*, 32, 7–41.
 重田康成・伊豆倉正隆・堤之恭, 2017. 北海道宗谷丘陵地域から産出した白亜紀末期・マーストリヒチアン前期アンモナイト。むかわ町穂別博物館研究報告, 32, 7–41.

(要旨)

北海道北部・宗谷丘陵地域の鬼志別川および猿骨川沿いに露出する蝦夷層群エタンパック層中部からマーストリヒチアン前期（白亜紀末期）のアンモナイトの産出を報告した。アンモナイトは8属9種で、*Neophylloceras hetonaiense* Matsumoto, *Tetragonites popetensis* Yabe, *T. terminus* Shigeta, *Gaudryceras izumiense* Matsumoto and Morozumi, *Anagaudryceras matsumotoi* Morozumi, *Pachydiscus* sp., *Diplomoceras* sp., *Nostoceras* sp., *Baculites regina* Obata and Matsumoto を含む。このアンモナイト群は、*Gaudryceras izumiense* と *Baculites regina* を含むことから、西南日本に分布する和泉層群の下部マーストリヒチアン階上部・*Gaudryceras izumiense* 帯の下部に対比できる。エタンパック層の最下部と最上部に挟在する凝灰岩中のジルコンの放射年代は、 72.6 ± 1.6 Ma と 70.6 ± 1.2 Ma であり、エタンパック層の年代がカンパニアン後期からマーストリヒチアン中期の最前期であることを示す。

Appendix 1. LA-ICP-MS analyzed data of zircons in the tuff samples collected from Loc. 3(OS001-025) and Loc. 10 (SY001-028) and calculated ages. Errors are 1 sigma. Pb_c and Pb* indicate the common and radiogenic portions, respectively.

Labels	$^{206}\text{Pb}_c$ ⁽¹⁾	U	Th	Th/U	$^{238}\text{U}/^{206}\text{Pb}^*$ ⁽¹⁾	$^{207}\text{Pb}^*/^{206}\text{Pb}^*$ ⁽¹⁾	$^{238}\text{U}/^{206}\text{Pb}^*$ age ⁽¹⁾	$^{238}\text{U}/^{206}\text{Pb}^*$ age ⁽²⁾
	(%)	(ppm)	(ppm)				(Ma)	(Ma)
OS001	0.72	475	280	0.60	91.17 ± 2.01	0.0369 ± 0.0069	70.3 ± 1.5	70.8 ± 1.5
OS002	0.00	611	717	1.20	93.82 ± 1.80	0.0555 ± 0.0036	68.3 ± 1.3	67.7 ± 1.3
OS003	0.77	931	1452	1.60	90.97 ± 2.16	0.0344 ± 0.0108	70.5 ± 1.7	71.0 ± 1.4
OS004	0.00	306	298	1.00	89.10 ± 2.49	0.0488 ± 0.0046	71.9 ± 2.0	71.8 ± 2.0
OS005	0.00	474	609	1.32	90.08 ± 2.19	0.0508 ± 0.0039	71.2 ± 1.7	70.9 ± 1.7
OS006	0.74	668	749	1.15	92.40 ± 2.09	0.0475 ± 0.0078	69.4 ± 1.6	69.4 ± 1.5
OS007	1.04	313	198	0.65	91.21 ± 2.63	0.0466 ± 0.0104	70.3 ± 2.0	70.4 ± 2.0
OS008	1.71	845	970	1.18	94.63 ± 1.94	0.0414 ± 0.0086	67.8 ± 1.4	68.3 ± 1.3
OS009	1.65	380	567	1.53	92.11 ± 2.56	0.0481 ± 0.0134	69.6 ± 1.9	69.6 ± 1.7
OS010	3.84	308	314	1.05	92.26 ± 2.66	0.0260 ± 0.0135	69.5 ± 2.0	71.4 ± 1.8
OS011	0.00	478	337	0.72	66.30 ± 1.22	0.0539 ± 0.0033	96.5 ± 1.8	95.8 ± 1.8
OS012	1.90	187	163	0.89	41.71 ± 1.01	0.0428 ± 0.0093	152.7 ± 3.6	153.9 ± 3.5
OS013	0.00	169	111	0.67	14.85 ± 0.25	0.0545 ± 0.0030	420.2 ± 6.9	420.2 ± 6.9
OS014	1.27	555	287	0.53	79.92 ± 1.61	0.0461 ± 0.0049	80.2 ± 1.6	80.3 ± 1.6
OS015	1.12	466	219	0.48	67.42 ± 1.57	0.0454 ± 0.0057	94.9 ± 2.2	95.2 ± 2.2
OS016	3.75	947	405	0.44	89.91 ± 1.67	0.0366 ± 0.0061	71.3 ± 1.3	72.3 ± 1.3
OS017	3.50	502	602	1.23	95.53 ± 2.31	0.0368 ± 0.0106	67.1 ± 1.6	68.0 ± 1.5
OS018	1.35	437	376	0.88	87.95 ± 2.09	0.0343 ± 0.0081	72.9 ± 1.7	73.9 ± 1.6
OS019	0.74	409	320	0.80	85.61 ± 1.83	0.0414 ± 0.0083	74.9 ± 1.6	75.4 ± 1.5
OS020	0.48	740	279	0.39	75.91 ± 1.29	0.0475 ± 0.0045	84.4 ± 1.4	84.4 ± 1.4
OS021	1.31	803	305	0.39	35.97 ± 0.53	0.0442 ± 0.0036	176.8 ± 2.6	178.0 ± 2.6
OS022	1.64	323	322	1.02	89.17 ± 2.38	0.0497 ± 0.0121	71.9 ± 1.9	71.7 ± 1.7
OS023	0.98	448	300	0.69	88.59 ± 2.09	0.0415 ± 0.0076	72.4 ± 1.7	72.9 ± 1.7
OS024	0.00	792	460	0.60	22.08 ± 0.29	0.0506 ± 0.0016	285.5 ± 3.7	285.5 ± 3.7
OS025	0.00	215	297	1.42	75.69 ± 2.46	0.0504 ± 0.0062	84.6 ± 2.7	84.3 ± 2.8
SY001	1.73	562	382	0.70	84.98 ± 1.85	0.0414 ± 0.0065	75.4 ± 1.6	76.0 ± 1.6
SY002	0.84	1697	587	0.36	71.16 ± 1.03	0.0490 ± 0.0029	90.0 ± 1.3	89.8 ± 1.3
SY003	0.00	732	194	0.27	2.46 ± 0.03	0.1471 ± 0.0012	2197.3 ± 22.6	2177.3 ± 22.6
SY004	3.28	812	736	0.93	74.38 ± 1.54	0.0605 ± 0.0073	86.1 ± 1.8	84.7 ± 1.7
SY005	0.00	302	111	0.38	68.63 ± 1.99	0.0805 ± 0.0055	93.3 ± 2.7	89.5 ± 2.7
SY006	0.00	474	650	1.41	66.61 ± 1.41	0.0757 ± 0.0038	96.1 ± 2.0	92.7 ± 2.0
SY007	0.00	387	295	0.78	71.24 ± 1.72	0.0641 ± 0.0044	89.9 ± 2.2	88.0 ± 2.2
SY008	0.29	429	294	0.70	30.01 ± 0.44	0.0537 ± 0.0041	211.3 ± 3.0	210.4 ± 2.9
SY009	1.83	119	28	0.24	47.54 ± 1.48	0.0482 ± 0.0079	134.2 ± 4.1	134.3 ± 4.1
SY010	1.68	589	392	0.68	88.03 ± 1.71	0.0364 ± 0.0067	72.8 ± 1.4	73.8 ± 1.4
SY011	1.73	194	125	0.66	85.77 ± 2.42	0.0526 ± 0.0108	74.7 ± 2.1	74.2 ± 2.0
SY012	1.59	576	546	0.97	91.66 ± 1.98	0.0410 ± 0.0075	69.9 ± 1.5	70.5 ± 1.5
SY013	2.05	516	166	0.33	78.87 ± 1.53	0.0486 ± 0.0056	81.2 ± 1.6	81.1 ± 1.5
SY014	3.43	150	97	0.67	88.60 ± 3.23	0.0326 ± 0.0149	72.3 ± 2.6	73.7 ± 2.6
SY015	0.00	292	135	0.47	81.38 ± 2.04	0.0485 ± 0.0049	78.7 ± 2.0	78.6 ± 2.0
SY016	0.00	576	326	0.58	89.43 ± 1.97	0.0490 ± 0.0035	71.7 ± 1.6	71.5 ± 1.6
SY017	0.57	231	133	0.59	2.72 ± 0.03	0.1470 ± 0.0026	2015.3 ± 20.2	1971.4 ± 20.1
SY018	0.60	397	338	0.87	72.89 ± 2.00	0.0793 ± 0.0109	87.8 ± 2.4	84.4 ± 2.2
SY019	1.82	277	144	0.53	70.15 ± 1.94	0.0657 ± 0.0100	91.3 ± 2.5	89.2 ± 2.4
SY020	1.14	278	233	0.86	75.07 ± 1.91	0.0465 ± 0.0097	85.3 ± 2.2	85.4 ± 2.0
SY021	0.23	732	280	0.39	58.05 ± 1.08	0.0556 ± 0.0042	110.1 ± 2.0	109.1 ± 2.0
SY022	0.00	598	276	0.47	89.08 ± 1.78	0.0584 ± 0.0056	72.0 ± 1.4	71.0 ± 1.4
SY023	0.63	327	215	0.67	84.20 ± 2.40	0.0498 ± 0.0090	76.1 ± 2.2	75.9 ± 2.1
SY024	0.34	747	195	0.27	58.91 ± 1.04	0.0494 ± 0.0041	108.5 ± 1.9	108.3 ± 1.9
SY025	1.77	420	238	0.58	77.54 ± 1.79	0.0488 ± 0.0083	82.6 ± 1.9	82.5 ± 1.8
SY026	0.00	242	142	0.60	73.39 ± 2.23	0.0734 ± 0.0069	87.2 ± 2.6	84.4 ± 2.7
SY027	0.16	807	276	0.35	73.93 ± 1.16	0.0420 ± 0.0038	86.6 ± 1.3	86.8 ± 1.3
SY028	0.00	303	107	0.36	58.79 ± 1.37	0.0545 ± 0.0046	108.7 ± 2.5	107.9 ± 2.6

Errors are 1-sigma; Pb_c and Pb* indicate the common and radiogenic portions, respectively.

(1) Common Pb corrected by assuming $^{206}\text{Pb}/^{238}\text{U}-^{208}\text{Pb}/^{232}\text{Th}$ age-concordance

(2) Common Pb corrected by assuming $^{206}\text{Pb}/^{238}\text{U}-^{207}\text{Pb}/^{235}\text{U}$ age-concordance

Appendix 2. List of ammonoids and inoceramids from the Etanpakk Formation of the Yezo Group exposed along the Onishibetsu and Sarukotsu rivers in the Soya Hill area, northernmost Hokkaido, northern Japan. Locality, sample (= concretion) and Hobetsu Museum (HMG) specimen numbers are shown. Suffix “p” of sample number means that the concretion was found as float. *Amat*, *Anagaudryceras matsumotoi*; *Breg*, *Baculites regina*; *Dipl*, *Diplomoceras* sp.; *Gizm*, *Gaudryceras izumiense*; *Nhet*, *Neophylloceras hetonaiense*; *Nost*, *Nostoceras* sp.; *Pachy*, *Pachydiscus* sp.; *Tpop*, *Tetragonites popetensis*; *Tter*, *Tetragonites terminus*.

locality	Sample	inoceramids	<i>Nhet</i>	<i>Amat</i>	<i>Gizm</i>	<i>Tter</i>	<i>Tpop</i>	<i>Pachy</i>	<i>Dipl</i>	<i>Nost</i>	<i>Breg</i>
	20150907-1					1720					
	20150907-3					1710					
	20150907-4										1742
	20150908-5	1768, 1769, 1770					1727				
1	20151007-1			1709							
1	20151007-2					1719					
	20150514-1p	1771					1728				
	20150514-3p						1729				
	20150610-1p		1702								
	20150907-4p	1772, 1773									
	20150514-9	1777			1712						
	20150514-10	1774, 1775, 1776									
	20150908-1	1779									
2	20150908-2	1778	1703, 1704		1711		1730				
	20150908-3					1731					
	20150514-7p	1780									
	20150514-8p	1781									
4	20150909-7	1755, 1756									
5	20150909-6	1757									
	20150514-11				1721	1732					1743
	20150514-12				1722						
6	20150909-2	1764, 1765, 1766									
	20150909-3	1762, 1763									
7	20150515-5						1739				
8	20150515-2p	1760, 1761	1705				1740				1744, 1745, 1746
	20150911-1p	1758, 1759									1747, 1748
9	20150513-1				1713						
	20150513-4	1767									
	20150515-14	1782				1733					1749, 1750, 1751, 1752, 1753
11	20150911-3p								1741		
	20160525-2p					1734					
	20160525-3p									1754	
	20150911-4				1714						
12	20150911-5	1783, 1784, 1785, 1786									
	20150912-1					1723	1735	1708			
	20151006-5					1724					
	20150515-15p				1715		1736				
13	20160523-4	1790				1725	1737				
	20150609-2				1716						
	20150710-1		1706								
	20150710-2						1738				
	20151008-2					1726					
14	20150515-8p	1787, 1788									
	20150515-9p	1789									
	20150515-10p		1707								
	20150515-11p				1718						
	20150609-1p				1717						
15	20150609-3	1791, 1792									